

*Original Paper*

# **A Possible Definition of a *Realistic* Physics Theory**

**Nicolas Gisin**

Group of Applied Physics, University of Geneva, 1211 Geneva 4, Switzerland

*E-mail:* [Nicolas.Gisin@unige.ch](mailto:Nicolas.Gisin@unige.ch)

Received: 9 October 2014 / Accepted: 24 November 2014 / Published: 1 December 2014

---

**Abstract:** A definition of a *Realistic* Physics Theory is proposed based on the idea that, at all time, the set of physical properties possessed (at that time) by a system should unequivocally determine the probabilities of outcomes of all possible measurements.

**Keywords:** quantum theory; realism; Bell inequality; nonlocality; propensity

---

## **1. Introduction**

Some of the physics literature of the last two decades abundantly mentions *realistic* theories, especially when discussing quantum physics, nonlocality and Bell inequalities (see e.g. [1–4]), or epistemic versus ontic interpretations of the quantum state [5,6]. On the one side, this is highly interesting as most physicists have some intuitive understanding of what "reality" is. However, on the other side, it is very disappointing, because the vast majority of this literature doesn't properly define *realistic* theories and thus obscures what could be an interesting debate [7].

The aim of this note is to propose a plausible definition of a *realistic theory*, section 2. It is not entirely new, but seems to have been forgotten. A good definition should be testable, however, the definition I propose is not testable in one single experiment. Nonetheless it defines whether a given physics theory is realistic or not. Hence, testing our best realistic physics theories amounts to testing whether we can maintain a realistic physical description of nature. Section 3 discusses nonlocality in the context of our definition.

## 2. Definition of *Realistic Physics Theories*

In the physics literature one finds essentially two characterizations of *realistic* theories:

1. All measurement outcomes are determined by the state of the physical system. In other words, at any time all physical quantities have their value somehow written in the physical system (these may change as time passes).
2. All measurement outcome probabilities are determined by the state of the physical system. In other words, at any time all physical quantities have the probabilities of all their possible values written in the physical system (again, these may change as time passes).

Note that both characterizations incorporate the idea that measurement outcomes (or their probabilities) are defined prior and independently of the observer, as one would expect for a realistic theory. Nevertheless, both characterizations are unsatisfactory. The first one identifies *realistic* with *deterministic*. Indeed, if measurement outcomes are *written in the system*, then, either randomness can only be due to our ignorance of the precise state of the system, i.e. randomness is not intrinsic but merely epistemic, or randomness is merely due to non-perfect measurements, i.e. randomness is technical as in classical physics. The second characterization is not better, as one can hardly see how a theory could not satisfy it, i.e. how the state of a physical system could not determine the probabilities of all measurement outcomes. Indeed, such "states" would make no prediction at all, they would not even predict the frequencies of measurement outcomes<sup>1</sup>. In brief, no physicist would call such "states", states of physical systems.

Consequently, the first characterization is too strong, as it imposes determinism, while the second characterization is too weak, as all scientific theories have to satisfy it. However, intuitively, both characterizations contain some truth. The state of physical system must determine the probabilities of all possible measurement outcomes, as suggested by the second characterization. And at least some physical properties<sup>2</sup> must be written in the physical system, as suggested by the first characterization (if not how would one even recognize the system?). This leads to the following definition:

---

<sup>1</sup> This is true whatever interpretation of probabilities one uses; indeed, whatever probabilities are, at the end of the day, probabilities are tested by comparing them with frequencies in long series of measurements.

<sup>2</sup> We distinguish physical quantities, like e.g. energy, and physical properties, like e.g. an energy eigenprojector. When testing a property the answer is binary, yes/no. A physical quantity corresponds to an entire set of compatible properties. In classical physics properties are represented by subsets of the phase-space, while in quantum theory properties are represented by projectors.

**Definition 1:** A theory is *realistic* if and only if, according to the mathematical structure of this theory, the collection of all physical quantities written in the system unambiguously determines the probabilities of all possible measurement outcomes.

In other words, the theory is such that, at all times, the collection of all properties possessed<sup>3</sup> by the system fully determines all relevant probabilities: the collection of physical quantities without uncertainty (or, more precisely, without indeterminacy) determines the probabilities of all undetermined physical quantities. Notice that a possessed property is similar to what EPR called an element of reality [8].

This definition captures both the idea that, at any time, some properties (physical quantities) are written in the system, as suggested by characterization number 1, and that what is written in the system determines all the relevant probabilities, as suggested by characterization number 2. When time passes, some properties possessed by the system pass away, while the system acquires new properties, i.e. the state of the system evolves.

Let us present our definition more rigorously (see also [9]). Our definition assumes that physics theories have some concept of *physical properties*, here denoted  $\pi$ , and of *probability measures*,  $\omega : \pi \rightarrow \omega(\pi) \in [0..1]$ . Furthermore, it is the theory that says which sets of properties  $\{\pi_j\}$  can be measured jointly in a single measurement (whose outcomes corresponding to the - thus compatible -  $\pi_j$ ). Probability measures must satisfy the following two natural conditions:

- i) If, according to the theory, a given measurement has several possibly mutually exclusive outcomes  $\pi_j$ , then, grouping several outcomes together<sup>4</sup>, denoted  $\Pi \equiv \{\pi_j\}$ , one has  $\omega(\Pi) = \sum_j \omega(\pi_j)$ .
- ii) If, for any measurement, one groups together all possible outcomes, then  $\omega(\mathbb{1}) = 1$ , where  $\mathbb{1}$  denotes the property corresponding to grouping together all outcomes.

A physics theory must also have some concept of *states*, here denoted  $\phi$ . Clearly, every state  $\phi$  should determine a unique probability distribution  $\omega_\phi$ . However, one should not identify states with probability measures. Indeed, if only the probabilities  $q_j$  of possible states  $\phi_j$ 's are known, then the probability measure reads  $\sum_j q_j \omega_{\phi_j}$ . Hence, not all probability measures correspond to a (pure) state. A priori, one could even have probability measures that do not correspond to any state, neither pure nor mixed.

If  $\omega_\phi(\pi) = 1$ , then, when the system is in state  $\phi$ , the outcome  $\pi$  is written in the physical system (here, a unit probability is identified with certainty).

<sup>3</sup> In this note I use equivalently the following terminologies: possessed property, actual property, property written in the system, i.e. properties such that the state of the system determines their yes/no values.

<sup>4</sup> The outcome  $\Pi$  corresponds to "one of the  $\pi_j$  happened".

We can now present our definition mathematically. A theory is realistic iff, according to the mathematical structure<sup>5</sup> of this theory, the following condition holds for all (pure) states  $\phi$  and all probability measures  $\omega$  that satisfy i) and ii):

If  $\{\pi|\omega(\pi) = 1\} = \{\pi|\omega_\phi(\pi) = 1\}$ , then  $\omega = \omega_\phi$ .

Hence, in realistic theories the state of a physical system can be identified with the collection of possessed properties. Indeed, this collection determines unequivocally all relevant probabilities.

This definition is in accordance with the idea that the state of a physical system is the collection of all "actual properties", i.e. of all physical quantities such that if one measures them, the outcome is certain or predetermined [10,11]. Accordingly, for any deterministic theory the state is merely a summary of all possible measurement results, as in classical mechanics and in some interpretations of quantum mechanics like Everett [12] and Bohm<sup>6</sup> [15]. For non-deterministic theories, the situation is more interesting [9]. For example, according to quantum theory, at any time  $t$ , there are physical quantities, represented in this theory by self-adjoint operators  $A_j$ , with well determined values  $a_j$ :  $A_j\Psi_t = a_j\Psi_t$ , where  $\Psi_t$  represents the quantum state at time  $t$ . Properties are represented by projectors, e.g. the eigenprojectors  $\pi_j^a$  of the operators  $A_j$  with spectral decompositions  $A_j = \sum_{a_j} a_j \cdot \pi_j^a$ .

Noteworthy, for any (normalized) pure state  $\Psi_t$ , the collection of all properties possessed by the system,  $\{\pi_j^a|A_j\Psi_t = a_j\Psi_t\}$ , determines a unique probability distribution  $\omega_{\Psi_t}$  such that  $\omega_{\Psi_t}(\pi_j^a) = 1$  for all outcomes  $\pi_j^a$  corresponding to a possessed property. This is nothing but Gleason's theorem (valid in all Hilbert spaces of dimension  $\geq 3$  [16]). As is well-known, this probability distribution reads  $\omega_{\Psi_t}(\pi_j^a) = \langle \Psi_t | \pi_j^a | \Psi_t \rangle$ .

According to Definition 1 both classical and quantum theories are realistic. Note that, in both theories, mixed states do not satisfy the proposed definition. This should be no surprise, as mixed states describe situations involving some ignorance, either about the precise state preparation or about a part of the global system which has been ignored (traced out, as physicists say).

The case of 2-dimensional quantum systems, so-called qubits, is interesting. Pure states can be labeled by vectors pointing on the unit sphere in ordinary 3-dimensional space,

<sup>5</sup> More precisely, the mathematical structure of the set of all properties, according to this theory.

<sup>6</sup> Notice that in Bohm's theory two apparata corresponding to the same self-adjoint operator and identical initial state-vector, but with different hidden positions of the particles making up the apparata may lead to different measurement outcomes (this is true even for position measurements [13]). Accordingly, in Bohm's theory there are many more properties than in quantum theory [14]. Consequently, the set of possessed properties in Bohm's theory defines not only the state-vector of the system, but also its hidden positions.

$\phi = |\vec{m}\rangle$ , and all nontrivial properties, (i.e. not equal to the identity  $\mathbb{1}$ ) can also be labeled by vectors on the unit sphere  $\pi_{\vec{p}} = |\vec{p}\rangle\langle\vec{p}|$ . Compatible (nontrivial) properties correspond to antipodal points on the sphere. According to quantum theory, every state  $|\vec{m}\rangle$  is associated to the following probability measure:  $\omega_{\vec{m}}(\pi_{\vec{p}}) = \langle\vec{m}|\pi_{\vec{p}}|\vec{m}\rangle = \frac{1+\vec{m}\cdot\vec{p}}{2}$ . Accordingly, for every state  $|\vec{m}\rangle$  there is a unique (nontrivial) actual property  $\pi_{\vec{m}}$  that corresponds to the same point on the sphere. However, since Gleason's theorem doesn't hold in dimension two, there are different probability measures with precisely the same set of actual properties. For example, the probability measure<sup>7</sup>  $\omega(\pi_{\vec{p}}) = \frac{1+(\vec{m}\cdot\vec{p})^3}{2}$  takes the value one on precisely the same set of properties as  $\omega_{\vec{m}}$ , but clearly differs from  $\omega_{\vec{m}}$ . Such an example is impossible for qutrits and all quantum systems described by Hilbert spaces of dimensions larger or equal to 3.

Consequently, quantum theory limited to one qubit is an example of a non-realistic theory. But, clearly, qubits don't exist per se, in some disembodied form, but only as subsystems of larger - actually infinite dimensional - systems; hence this is a nice example, but not a compelling one against fundamental theories being realistic.

Let me conclude this section by admitting that I would be greatly surprised, even shocked, if any future fundamental theory in physics would be non-realistic according to the proposed definition<sup>8</sup>. But logically this is certainly possible.

### 3. Nonlocality

In the context of quantum nonlocality, some physicists conclude that no theory compatible with the violation of Bell inequalities can be realistic. Indeed, some claim that the negation of "local realism" leaves us with a choice to either give up locality or realism. However, as long as the word "realistic" is not defined, one can't argue against (nor in favor) of such a statement [17]. According to Definition 1, such a statement is clearly wrong, as quantum theory predicts the violation of some Bell inequalities and is realistic<sup>9</sup>. However, in order for a realistic theory to predict the violation of some Bell

<sup>7</sup> Note that  $\omega(\pi_{\vec{p}}) + \omega(\pi_{-\vec{p}}) = \omega(\mathbb{1}) = 1$ , as it should.

<sup>8</sup> For me, realism is merely the existence of a world outside me and independent of me. A priori, this world could be described as well by classical physics as by quantum theory or by any future theory. Indeed, the existence of an outside world doesn't impose any restriction on how this world is (the main and only point is that it exists). This outside world could be deterministic or random, local or nonlocal, entirely within space-time or not, fully understandable by our intelligence or not. Hence, physics will never be able to deny its existence. However, let me confess an act of faith: I believe that by studying physics I open a small skylight on this external world. What I can see is only a tiny part of it, but already this tiny part is absolutely fascinating: enlarging this small window is the great Enterprise of Science. And let me add that if one assumes realism, then it is all the most natural to look for realistic theories and to be satisfied only by realistic theories.

<sup>9</sup> One may ask whether a non-realistic local theory is still an option. However, as long as a theory makes

inequalities, the theory must incorporate some form of nonlocality. Definition 1 of a *realistic theory* rests on the concept of *possessed properties*, or equivalently of *physical quantities without indeterminacy*. Thus, if the system is delocalized, like a delocalized photon in two (or more) optical modes or like two (or more) entangled particles located at a distance from each other, then some of the physical quantities without indeterminacy are also delocalized. Note that in order to measure such delocalized physical quantities the measurement apparatus itself must be delocalized or must consist of several parts. It is certainly highly counter-intuitive that some properties written (at some point in time) in a physical system can be delocalized, but that doesn't make them less real. For example, if two systems have their spin  $\frac{1}{2}$  entangled as in the singlet state, then their global spin is zero although this global spin is delocalized; one way to measure it uses an apparatus with two pieces, one near each particle, each measuring the spin  $\frac{1}{2}$  along the same direction; the pre-determined outcome is that the two local results are opposite. In this example, the possessed property *global spin zero* is written in the 2-particle system, hence it is written out there in space, but in a nonlocal fashion.

#### 4. Conclusion

I proposed a possibly definition of realistic theories which avoids the mere reduction to determinism and avoids the risk of tautology. This definition is closely related to the concept of state as the collection of all actual properties, or equivalently as the collection of all elements of reality. According to it, both classical and quantum theories are realistic. Consequently, one can't avoid nonlocality by merely rejecting realistic theories. Still, non-realistic theories exist mathematically and one may wonder whether any future fundamental physics theory may be non-realistic.

Other definitions of realistic physics theories are possible [18]. It is an interesting research program to compare them and to investigate the status of Bell-nonlocality in each of them.

#### Acknowledgment

It is a pleasure to thank Tomer Barnea, Jeff Bub, Michael Esfeld, Yeong-Cherng Liang, Gilles Puetz, Terry Rudolf and Valerio Scarani for helpful comments. Financial support by the Swiss SNF grant 200021\_149109 is gratefully acknowledged.

---

predictions, i.e. is a scientific theory, it determines probabilities, hence allows one to define Bell-locality as the factorization of probabilities corresponding to distant events. Accordingly, Bell inequality can be derived in all theories that make predictions: no local scientific theory is compatible with the violation of Bell's inequality, independently of whether the scientific theory is realistic or not.

## References

1. S. Gröblacher et al., [An experimental test of non-local realism](#), Nature **446**, 871 (2007); T. Paterek et al. [Experimental Test of Nonlocal Realistic Theories Without the Rotational Symmetry Assumption](#), Phys. Rev. Lett. **99**, 210406 (2007).
2. M. Giustina et al., [Bell violation using entangled photons without the fair-sampling assumption](#), Nature **497**, 227 (2013).
3. J. Hofmann et al, [Heralded Entanglement Between Widely Separated Atoms](#), Science **337**, 72-75 (2012).
4. B. G. Christensen et al., [Detection-Loophole-Free Test of Quantum Nonlocality, and Applications](#), Phys. Rev. Lett. **111**, 130406 (2013).
5. N. Harrigan and R. W. Spekkens, [Einstein, incompleteness, and the epistemic view of quantum states](#), Foundations of Physics **40**, 125 (2010).
6. M. F. Pusey, J. Barrett and T. Rudolph, [On the reality of the quantum state](#), Nature Phys. **8**, 476 (2012)
7. N. Gisin, [Quantum nonlocality: how does nature do it?](#), Science, **326**, 1357-8 (2009).
8. A. Einstein, B. Podolsky and N. Rosen, [Can Quantum-Mechanical Description of Physical Reality Be Considered Complete?](#), Phys. Rev. **47**, 777 (1935).
9. N. Gisin, [Propensities in a non Deterministic Physics](#), Synthese **89**, 287-297 (1991).
10. C. Piron, [Foundations of Quantum Physics](#), W.A. Benjamin, Inc., 1976.
11. C. Piron, [A First Lecture on Quantum Mechanics](#), in [Quantum Mechanics, a Half Century Later](#), Editors J. Leite Lopez and M. Paty, Reidel Pub. Co, pp 69-87, 1977.
12. H. Everett III, ["Relative State" Formulation of Quantum Mechanics](#), Rev. Mod. Phys. **29**, 454 (1957).
13. J. Kiukas and R.F. Werner, [Maximal violation of Bell inequalities by position measurements](#), J. Math. Phys. **51**, 072105 (2010).
14. M. Daumer, D. Dürr, S. Goldstein and N. Zanghi, [Naive Realism about Operators](#), in [Probability, Dynamics and Causality](#), pp 239-257, Springer, 1997; [arXiv/quant-ph/9601013](#).
15. D. Bohm, [A suggested Interpretation of the Quantum Theory in Terms of "Hidden" Variable](#), Phys. Rev., **85**, 166-193 (1952); D. Bohm and B. J. Hilley, [The Undivided Universe](#), New York: Routledge, 1993.
16. A. Peres, [Quantum Theory: Concepts and Methods](#), Kluwer 1993.
17. N. Gisin, [Non-realism: deep thought or a soft option?](#), Found. Phys. **42**:80Öy5, 2012.
18. J. Bub and R. Clifton, [A uniqueness Theorem for "No Collapse" Interpretation of Quantum Mechanics](#), Stud. Hist. Phil. Mod. Phys. **27**, 181-219 (1996).